
Special Project Palliative Care

Medical Directive

A Primary Care Paramedic may provide the treatment, transport and/or referral prescribed in this Medical Directive for registered patients if authorized.

These registered palliative care patients require a different approach to assessment and treatment that reflects their goals of care. Therefore paramedics, for this defined patient population, should prioritize patient comfort and are not required to follow the described regimen of strict vital signs, cardiac monitoring and transport as directed in the Basic Life Support Patient Care Standard (BLS PCS). If patient transport is initiated however, paramedics should proceed with usual care per the BLS PCS. If a paramedic determines that the patient would benefit from any other medical directives in the Advanced Life Support (ALS) PCS that is not included in this special project medical directive, a patch to a base hospital physician (BHP) is necessary.

DYSPNEA

INDICATIONS

Patient registered in palliative care program

And

Uncontrolled dyspnea with suspected bronchoconstriction

CLINICAL CONSIDERATIONS

- ▶ Salbutamol should only be used in patients whose dyspnea is accompanied by wheezing or a history of response to bronchodilators.

CONDITIONS

Salbutamol

AGE: ≥18

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: For Dyspnea
with suspected
bronchoconstriction
only

CONTRAINDICATIONS

Salbutamol

Allergy to salbutamol

TREATMENT



Patient • Drug • Dose • Route • Time.

Consider **Salbutamol**

	Route <i>MDI*</i>	Route <i>NEB</i>
<i>Dose</i>	Up to 800 mcg (8 puffs)	5 mg
<i>Max. dose</i>	800 mcg	5mg
<i>Dosing interval</i>	5-15 min prn	5-15 min prn
<i>Max. # of doses</i>	3	3

*1 puff – 100 mcg

HALLUCINATIONS OR AGITATION

INDICATIONS

Patient registered in palliative care program

And

Increasing agitation or suspected new or increased hallucinations

CLINICAL CONSIDERATIONS

CONDITIONS

Haloperidol

AGE: ≥ 18

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

CONTRAINDICATIONS

Haloperidol

Allergy to haloperidol

Known Parkinson's or Lewy
Body Dementia

Neuroleptic Malignant
Syndrome

TREATMENT



Patient • Drug • Dose • Route • Time.

Consider **Haloperidol**

	Route
	SC
<i>Dose</i>	0.5-1 mg
<i>Max. single dose</i>	1 mg
<i>Dosing interval</i>	30 min
<i>Max. # of doses</i>	2

NAUSEA OR VOMITING

INDICATIONS

Patient registered in palliative care program

And

Nausea and/or vomiting

CLINICAL CONSIDERATIONS

- ▶ Dimenhydrinate is rarely used in the palliative care population as it can cause delirium, increase drowsiness, and does not target the appropriate receptors to control the nausea in most patients. It should only be used in patients with contraindications to haloperidol where ondansetron cannot be used and should be started at low doses.

CONDITIONS

Haloperidol	Ondansetron	Dimenhydrinate
AGE: ≥18	AGE: ≥18	AGE: ≥18
LOA: N/A	LOA: N/A	LOA: N/A
HR: N/A	HR: N/A	HR: N/A
RR: N/A	RR: N/A	RR: N/A
SBP: N/A	SBP: N/A	SBP: N/A
Other: N/A	Other: Contraindication to Haloperidol	Other: Contraindication to Haloperidol

CONTRAINDICATIONS

Haloperidol	Ondansetron
Allergy to haloperidol Known Parkinson's or Lewy Body Dementia Neuroleptic Malignant Syndrome	Allergy to ondansetron

Dimenhydrinate
Allergy to dimenhydrinate or other antihistamines Overdose on antihistamines or anticholinergics or tricyclic antidepressants

TREATMENT



Patient • Drug • Dose • Route • Time.

Consider **Haloperidol**

	Route
	SC
<i>Dose</i>	0.5-1 mg
<i>Max. single dose</i>	1 mg
<i>Dosing interval</i>	30 min
<i>Max. # of doses</i>	2

Consider **Ondansetron**

	Route
	<i>PO/SC</i>
<i>Dose</i>	4 mg
<i>Max. single dose</i>	4 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

Consider **Dimenhydrinate**

	Route
	<i>SC</i>
<i>Dose</i>	25-50 mg
<i>Max. single dose</i>	50 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

TERMINAL CONGESTED BREATHING

INDICATIONS

Patient registered in palliative care program

And

Congested/loud/rattling breathing in patients near the end of life

CLINICAL CONSIDERATIONS

- ▶ Patient repositioning and gentle turning of the head to the side can be done instead of medication however suction of the oropharynx is not appropriate as it will likely cause discomfort and a gag reflex.

CONDITIONS

Glycopyrrolate or Atropine

AGE: ≥ 18

LOA: N/A

HR: N/A

RR: N/A

SBP: N/A

Other: N/A

CONTRAINDICATIONS

Glycopyrrolate

Allergy to glycopyrrolate

Atropine

Allergy to atropine

TREATMENT



Patient • Drug • Dose • Route • Time.

Consider **Glycopyrrolate or Atropine**

	Route
	<i>SC</i>
<i>Dose</i>	0.4 mg
<i>Max. single dose</i>	0.4 mg
<i>Dosing interval</i>	N/A
<i>Max. # of doses</i>	1

TREAT AND REFER

INDICATIONS

Patient registered in palliative care program

And

Symptoms improved to patient's/Substitute Decision Maker's (SDM) satisfaction

And

After informed discussion patient/SDM preference to remain at home

CLINICAL CONSIDERATIONS

- ▶ A period of observation is recommended after the administration of any medication if the patient is not transported to ensure adequate response and no unexpected immediate adverse effects
 - ▶ Transport should be considered if there is strong suspicion of reversible causes including but not limited to:
 - Complete bowel obstruction with no prior history of same
 - New Spinal Cord Compression
 - New Superior Vena Cava (SVC) Obstruction
 - Airway obstruction
 - Suspected new pathologic fracture
 - ▶ If patients do not meet the treat and refer conditions, paramedics should consider consulting BHP, follow the patient refusal standard and document appropriately.
-

CONDITIONS

Age \geq 18

Valid DNR Confirmation Form

Patient registered in Paramedic Palliative Care Program

CONTRAINDICATIONS

Concerns of patient abuse or neglect

Patient and SDM cannot demonstrate decision-making capacity based on the Aid to Capacity Evaluation Tool

Uncontrolled or new seizures

TREATMENT

Paramedics may treat patients according to this medical directive and, in collaboration with the patient/SDM, honour wishes to remain at home (treat and refer). Paramedics will notify the patient's palliative care team.