

## Centre for Prehospital Care (CPC) **STANDARDS**

CATEGORY: Program Specific REVISION DATE:

**ISSUE DATE:** March 2023

TITLE: Discontinuing Cardiac Monitor Use Page 1 of 2

Doorwood Owner	Name	
Document Owner:	Name:	
Regional Manager	Nicole Sykes	
Update Schedule: Annually		
Stakeholder Consultation and Review:	Date:	
Centre for Prehospital Care Quality of Care	March 23, 2023	
Committee		
Centre for Prehospital Care Program Committee	April 11, 2023	
·	·	
Approval:	Date:	
Nicole Sykes, Chair, CPC Program Council	April 19, 2023	
Electronic Signature	·	

## **PURPOSE**

To provide guidance for when a paramedic may discontinue cardiac monitor use. The BLS PCS identifies cardiac monitoring as a necessary assessment method for certain patients. However, many patients who have had cardiac monitoring will not require it on a continued or ongoing basis.

## **STANDARDS**

- As per the ALS PCS and BLS PCS:
  Paramedics will apply the cardiac monitor to the patient.
- Paramedics will keep the cardiac monitor applied to the patient while determining final patient disposition. If the patient is transported, paramedics will continue the cardiac monitor while en route to the receiving facility.
- Where a cardiac monitor has been applied by paramedics, the cardiac monitor may be removed if:
  - Patient's vitals are consistently within normal parameters as outlined in the ALS PCS Preamble, Age and Vital Signs section or explainable by or not pertinent to the presenting complaint based on paramedic judgement;
  - The patient has not complained of ischemic type chest pain or palpitations at any point;
  - There is no incident history of loss of consciousness (of any duration);
  - Patient shows no electrical cardiac activity that is abnormal or inconsistent with previous medical history (e.g. rate or rhythm);
  - The patient exhibits no signs of respiratory distress;
  - There is no incident history of cerebrovascular accident (CVA) or transient ischemic attack (TIA);
  - is neither confirmed nor suspected in the incident history of ingesting drugs, medications, or other substances that may lead to toxicity;
  - No other ALS PCS interventions are anticipated;
  - The patient has had no further symptoms requiring medication and peak onset of any administered medication has been achieved; and
  - If an IV is in place, it is TKVO (30-60 ml/hr adult) or a saline lock is applied.

CATEGORY: i.e. Program Specific Page 2 of 2

TITLE: NAME OF DOCUMENT

• If the conditions above are met, the paramedic may remove the cardiac monitor and document on the eACR with the appropriate code, that the cardiac monitor has been discontinued and the reason. After removal of the cardiac monitor, the patient remains the responsibility of the paramedic until patient disposition is finalized and/or if transport has occurred, there is a transfer of care to appropriate staff.

• Following discontinuation of cardiac monitoring, paramedics who continue to provide care to the patient shall provide appropriate reassessment of the patient as indicated by Patient Care Standards until care is formally transferred to appropriate staff. Reassessment includes measurement of vital signs at intervals suitable to the patient's condition.

Date/Time	Code	Procedure	Results
Jan. 25, 2023 11:11	314	Cardiac Monitor	Monitor disconnected and the patient
		Disconnect	remained with Paramedic 181XX

## **References and Related Documents**

Advanced Life Support Patient Care Standards (ALS PCS) Basic Life Support Patient Care Standards (BLS PCS)